



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

LYCOPodium COMPLANATUM VAR. **FLABELLIFORME** WITH SEVEN SPIKES.—In recently working over some undistributed material in the Pennsylvania herbarium at the Carnegie Museum I found a specimen of *Lycopodium complanatum* var. *flabelliforme* Fernald, apparently typical of that variety excepting that the fruiting stem carried two peduncles, each subtending seven well developed spikes averaging about two centimeters long. The distances between successive forks, as well as the immediate pedicels of the spikes, range from about 4 to 8 mm., but at the second forking of the peduncle one of the branches remains undivided and constitutes thus a pedicel for its spike about 11 mm. long. The specimen was collected by O. P. Medsger, Dec. 27, 1899, along Jacobs Creek, which flows in a rather wild valley in an almost mountainous region of southwestern Pennsylvania.—O. E. JENNINGS *Pittsburgh, Pa.*

A NEW STATION FOR CYRTOMIUM FALCATUM AND PTERIS LONGIFOLIA IN ALABAMA.—Mr. W. C. Dukes reported in the Fern Bulletin for July, Vol. XV, No. 3, of having located a station for *Cyrtomium falcatum* (*Aspidium falcatum*) near Prattville, in Autauga County, Alabama.

Last year I noticed a fern growing along the street near the Ashland Place, on the north side of Spring Hill road, just west of the city limits of Mobile. This year I examined it more closely and found it to be a good sized plant of *Cyrtomium falcatum*. As it was growing in the corner of a yard of a dwelling house, I called at the house and asked the occupants if they had set the fern there. On their replying in the negative, I asked how long it had been growing there, I was told it was there four years previous when they moved into the house.

It appears to be a very old plant for the rhizome is large and covered with rough scales, and has grown out in two directions, the ends being about eight inches apart. One end was dead, but the other though still alive was being used badly by chickens that had been scratching at it. It was growing in a rather dry place near the end of a brick wall under *Crepe Myrtle* (*Lagerstroemia indica*, L.) bushes. There might have been more plants but for the chickens.

I found a dozen or fifteen plants of *Pteris longifolia* L. growing on the brick wall of an old building on Congress street, near Davis Avenue in the city of Mobile. There are growing with it *Pteris serrulata*, L. and *Dryopteris patens*, (Sw.) Kuntze and several kinds of weeds. The wall is covered with moss and is shaded by a tree. I have observed it growing there for two years. It may be found in other parts of the city but this the only place I have seen it growing. This fern is not reported in Mohr's Plant Life of Alabama, and as far as I know this is the first time it has been reported from the state. —E. W. GRAVES, *Spring Hill, Ala.*

ON THE VIABILITY OF CERTAIN FERN SPORES.—The spores of some of our common wild ferns germinate only a short time after they have reached maturity as, for example, those of the *Osmunda* species which remain viable for only a few days. In other species as *Pteris aquilina* L. the spores are known to retain their power to germinate for two years.

Beginning with the summer of 1910, the writer collected in the vicinity of Madison fronds of a number of species of ferns. The spores of all the species which germinated a short time after they were collected were kept in packets in the botanical laboratory. The spores of the different species were sown again October 4,